arXiv:1502.05527v1 [math.FA] 19 Feb 2015

COMPACT EMBEDDING DERIVATIVES OF HARDY SPACES INTO LEBESGUE SPACES

JOSÉ ÁNGEL PELÁEZ

Abstract. We characterize the positive Borel measures such that the differentiation operator of order $n \in \mathbb{N} \cup \{0\}$ is compact from the Hardy space H^p into $L^q(\mu)$, $0 < p, q < \infty$.

1. Introduction

Let $\mathbb{D}$ denote the open unit disk of the complex plane and let $\mathbb{T}$ denote the unit circle. Also, let H^p, $0 < p < \infty$ denote the standard Hardy space of analytic functions in $\mathbb{D}$.

The aim of this paper is to characterize the positive Borel measures μ on the unit disc $\mathbb{D}$ such that the differentiation operator $D^{(n)}(f) = f^{(n)}$ is compact from H^p into $L^q(\mu)$, $n \in \mathbb{N} \cup \{0\}$ and $0 < p, q < \infty$.

The analogous problem for the standard Bergman spaces A^p_α has been solved [9, 11, 14]. The formula $||f||_{A^p_\alpha} \asymp \sum_{j=0}^{n-1} |f^{(j)}(0)| + ||f^{(n)}||_{A^p_{\alpha+np}}$ implies that for these spaces the question of when the differentiation operator $D^{(n)}$ from A^p_α into $L^q(\mu)$ is bounded or compact, can be answered once the case $n = 0$ is solved. However, this method does not work for Hardy spaces, because such a Littlewood-Paley formula does not exist for $p \neq 2$. Nevertheless, an equivalent H^p-norm in terms of the nth derivative can be given by using the square functions

$$S_{\sigma,n} f(\zeta) = \left(\int_{\Gamma_\sigma(\zeta)} |f^{(n)}(z)|^2 (1 - |z|)^{2n-2} \, dA(z) \right)^{1/2}$$

where $\Gamma_\sigma(\zeta) = \{z \in \mathbb{D} : |\arg \zeta - \arg z| < \sigma\,(1 - |z|)\}$ denotes the Stolz angle (lens type region) with vertex at $\zeta \in \mathbb{T}$ and aperture $\sigma > 0$. Precisely, for $0 < p < \infty$ [1, 8]

$$\|f\|^p_{H^p} \asymp \sum_{j=0}^{n-1} |f^{(j)}(0)|^p + \int_{\mathbb{T}} S^p_{\sigma,n} f(\zeta) \, dm(\zeta) \,. \tag{1.1}$$

Date: February 20, 2015.

2010 *Mathematics Subject Classification.* 30H10.

Key words and phrases. Hardy spaces, tent spaces, Carleson measures, differentiation operator, compact operators.

The author was supported in part by the Ramón y Cajal program of MICINN (Spain), Ministerio de Educación y Ciencia, Spain, (MTM2011-25502), from La Junta de Andalucía, (FQM210) and (P09-FQM-4468).

Here and throughout in what follows m denotes the arclength measure on $\mathbb{T}$. In view of (1.1) and the H^p characterization through the non-tangential maximal function, it is natural that Luecking [10, 11] employed tent spaces to describe the positive Borel measures such that $D^{(n)}: H^p \to L^q(\mu)$ is bounded. It is worth noticing that Coifman, Meyer and Stein [6] introduced the theory of tent spaces in an harmonic analysis context. It was further extended by Cohn and Verbitsky [4] and has become a very useful tool in operator theory on Hardy spaces [4, 5].

Let us now recall some definitions that will enable to state the solution to the primary question of this paper. Let us write $\Gamma_{1/2}(\zeta) = \Gamma(\zeta)$ for short, and for each $z \in \mathbb{D}$ let be $I(z) = \{\zeta \in \mathbb{T} : z \in \Gamma(\zeta)\}$ the related interval. The Carleson square $S(I)$ based on an interval $I \subset \mathbb{T}$ is the set $S(I) = \{re^{it} \in \mathbb{D} : e^{it} \in I, \, 1 - |I| \le r < 1\}$, where $|E|$ denotes the Lebesgue measure of $E \subset \mathbb{T}$.

If $|I| < \frac{\pi}{4}$ the tent $T(I)$ is the open subset of $\mathbb{D}$ bounded by the arc $I \subset \mathbb{T}$ and two straight lines through the endpoints of I forming with I an angle of $\frac{\pi}{4}$. If $|I| \ge \frac{\pi}{4}$, we set $T(I) = \cup_{J \subset I, |J| < 1} T(J) \cup \{0\}$. For each $a \in \mathbb{D}$, let be $S(a) = S(I(a))$ and $T(a) = T(I(a))$.

For $0 < q < \infty$ and a positive Borel measure ν on $\mathbb{D}$, finite on compact sets, denote $A^q_{q,\nu}(f)(\zeta) = \int_{\Gamma(\zeta)} |f(z)|^q \, d\nu(z)$ and $A_{\infty,\nu}(f)(\zeta) = \nu\text{-ess}\sup_{z \in \Gamma(\zeta)} |f(z)|$. For $0 < p < \infty$, $0 < q \le \infty$ the tent space $T^p_q(\nu)$ consists of the ν-equivalence classes of ν-measurable functions f such that $\|f\|_{T^p_q(\nu)} = \|A_{q,\nu}(f)\|_{L^p(\mathbb{T},m)}$ is finite. For $0 < q < \infty$, define

$$C^q_{q,\nu}(f)(\zeta) = \sup_{a \in \Gamma(\zeta)} \frac{1}{|I(a)|} \int_{T(a)} |f(z)|^q (1 - |z|) \, d\nu(z), \quad \zeta \in \mathbb{T}.$$

A quasi-norm in the tent space $T^\infty_q(\nu)$ is defined by $\|f\|_{T^\infty_q(\nu)} = \|C_{q,\nu}(f)\|_{L^\infty(\mathbb{T},m)}$.

The following result gives a description of the dual of $T^p_q(\nu)$ [6, 10]. We also refer to [13, Theorem 4] where an analogue was proved for a family of weighted tent spaces on the unit disc.

Theorem A. *Let $1 \le p, q < \infty$ with $p + q \ne 2$ and let ν be a positive Borel measure on $\mathbb{D}$, finite on compact sets of $\mathbb{D}$. Then the dual of $T^p_q(\nu)$ can be identified with $T^{p'}_{q'}(\nu)$ (up to an equivalence of norms) under the pairing*

$$\langle f, g \rangle_{T^2_2(\nu)} = \int_{\mathbb{D}} f(z) \overline{g(z)} (1 - |z|) \, d\nu(z).$$

For the sake of completeness, and because it is a key to describe those positive Borel measures such that $D^{(n)}: H^p \to L^q(\mu)$ is compact, we shall prove in Section 2 that each $g \in T^{p'}_{q'}(\nu)$ induces a bounded linear functional on $T^p_q(\nu)$. In the proof for $p = 1$, a stopping time involving $A_{q,\nu}(f)$ and $C_{q,\nu}(f)$ is a fundamental step. Let $\Delta(a, r)$ and $D(a, r)$ respectively denote the pseudohyperbolic and Euclidean discs of center a and radius r. Our main result is the following.

Theorem 1. *Let $0 < p, q < \infty$, $n \in \mathbb{N}$ and μ be a positive Borel measure on $\mathbb{D}$. Further, let $dh(z) = dA(z)/(1 - |z|^2)^2$ denote the hyperbolic measure.*

(a) *If* $p \geq q$, $D^{(n)} : H^p \to L^q(\mu)$ *is compact if and only if, for any fixed* $r \in (0,1)$, *the function*

$$\Phi_\mu(z) = \frac{\mu(\Delta(z,r))}{(1-|z|)^{1+qn}}, \quad z \in \mathbb{D},$$

satisfies that

  (i) $\Phi_\mu \in T^{\frac{p}{p-q}}_{\frac{2}{2-q}}(h)$, *if* $q < \min\{2, p\}$;

  (ii) $\lim_{|a|\to 1^-} \frac{1}{|I(a)|} \int_{T(a)} |\Phi_\mu(z)|^{\frac{2}{2-p}} \frac{dA(z)}{1-|z|} = 0$, *if* $q = p < 2$;

  (iii) $\lim_{R\to 1^-} \int_{\mathbb{T}} \left(\sup_{z \in \Gamma(\zeta) \setminus \overline{D(0,R)}} \Phi_\mu(z) \right)^{\frac{p}{p-q}} dm(\zeta) = 0$, *if* $2 \leq q < p$.

(b) *If either* $q > p$ *or* $2 \leq q = p$, *the following conditions are equivalent:*

  (i) $D^{(n)} : H^p \to L^q(\mu)$ *is compact;*

  (ii) $\lim_{|z|\to 1^-} \frac{\mu(S(z))}{(1-|z|)^{\frac{q}{p}+nq}} = 0$;

  (iii) $\lim_{|z|\to 1^-} \frac{\mu(\Delta(z,r))}{(1-|z|)^{\frac{q}{p}+nq}} = 0$ *for any fixed* $r \in (0,1)$.

As for $n = 0$, $I_d : H^p \to L^q(\mu)$ is compact if and only if $\lim_{|z|\to 1^-} \frac{\mu(S(z))}{(1-|z|)^{\frac{q}{p}}} = 0$, whenever $0 < p \leq q < \infty$ [3]. In the previous condition $\mu(S(z))$ may be replaced by $\mu(\Delta(z,r))$ if $p < q$. In the triangular case $0 < q < p < \infty$, $I_d : H^p \to L^q(\mu)$ is bounded if and only the function $B_\mu(\zeta) = \int_{\Gamma(\zeta)} \frac{d\mu(z)}{1-|z|}$ belongs to $L^{\frac{p}{p-q}}(\mathbb{T}, m)$ [10]. For this last range of values it is probably known, at least to experts working on the field, that $I_d : H^p \to L^q(\mu)$ is compact if and only it is bounded. Since we were not able to find a proof in the existing literature, we include a proof here.

Theorem 2. *Let* $0 < q < p < \infty$ *and let* μ *be a positive Borel measure on* $\mathbb{D}$. *Then the following conditions are equivalent:*

  (i) $I_d : H^p \to L^q(\mu)$ *is compact;*

  (ii) $I_d : H^p \to L^q(\mu)$ *is bounded;*

  (iii) *The function* $B_\mu(\zeta) = \int_{\Gamma(\zeta)} \frac{d\mu(z)}{1-|z|}$ *belongs to* $L^{\frac{p}{p-q}}(\mathbb{T}, m)$.

Throughout the paper $\frac{1}{p} + \frac{1}{p'} = 1$. Further, the letter $C = C(\cdot)$ will denote an absolute constant whose value depends on the parameters indicated in the parenthesis, and may change from one occurrence to another. We will use the notation $a \lesssim b$ if there exists a constant $C = C(\cdot) > 0$ such that $a \leq Cb$, and $a \gtrsim b$ is understood in an analogous manner. In particular, if $a \lesssim b$ and $a \gtrsim b$, then we will write $a \asymp b$.

2. Preliminary results

Proposition B. *Let* $1 \leq p, q < \infty$ *with* $p + q \neq 2$ *and let* ν *be a positive Borel measure on* $\mathbb{D}$, *finite on compact sets of* $\mathbb{D}$. *Then, there exists a positive constant* C *such that*

$$\left| \langle f, g \rangle_{T_2^2(\nu)} \right| \leq C ||f||_{T_q^p(\nu)} \|g\|_{T_{q'}^{p'}(\nu)}$$

for any $f \in T^p_q(\nu)$ *and* $g \in T^{p'}_{q'}(\nu)$.

Proof. If $1 < p, q < \infty$, then Fubini's theorem and two applications of Hölder's inequality give

$$|\langle f, g\rangle_{T^2_2(\nu)}| \le \int_{\mathbb{T}} A_{q,\nu}(f)(\zeta) A_{q',\nu}(g)(\zeta)\, dm(\zeta) = \|f\|_{T^p_q(\nu)} \|g\|_{T^{p'}_{q'}(\nu)}. \tag{2.1}$$

If $q = 1$ and $1 < p < \infty$, then Hölder's inequality yields

$$|\langle f, g\rangle_{T^2_2(\nu)}| \le \int_{\mathbb{T}} A_{1,\nu}(f)(\zeta) A_{\infty,\nu}(g)(\zeta)\, dm(\zeta) = \|f\|_{T^p_1(\nu)} \|g\|_{T^{p'}_{\infty}(\nu)}. \tag{2.2}$$

Let now $p = 1$ and $1 < q < \infty$. For $\zeta \in \mathbb{T}$ and $0 \le h \le \infty$, let

$$\Gamma^h(\zeta) = \Gamma(\zeta) \setminus \overline{D\left(0, \frac{1}{1+h}\right)} = \left\{ z \in \mathbb{D} : |\arg z - \arg \zeta| < \frac{1 - |z|}{2} < \frac{h}{2(1+h)} \right\}$$

and

$$A^{q'}_{q',\nu}(g|h)(\zeta) = \int_{\Gamma^h(\zeta)} |g(z)|^{q'}\, d\nu(z), \quad \zeta \in \mathbb{T}.$$

For every $g \in T^{\infty}_{q'}(\nu)$ and $\zeta \in \mathbb{T}$, define the stopping time by

$$h(\zeta) = \sup\left\{ h : A_{q',\nu}(g|h)(\zeta) \le C_1 C_{q',\nu}(g)(\zeta) \right\},$$

where $C_1 > 0$ is a large constant to be determined later. Assume for a moment that there exists a constant $C_2 > 0$ such that

$$\int_{\mathbb{D}} k(z)(1 - |z|)\, d\nu(z) \le C_2 \int_{\mathbb{T}} \left(\int_{\Gamma^{h(\zeta)}(\zeta)} k(z)\, d\nu(z) \right) dm(\zeta) \tag{2.3}$$

for all ν-measurable non-negative functions k. Then, applying Hölder's inequality

$$\begin{aligned} |\langle f, g\rangle_{T^2_2(\nu)}| &\le C_2 \int_{\mathbb{T}} \left(\int_{\Gamma^{h(\zeta)}(\zeta)} |f(z)||g(z)|\, d\nu(z) \right) dm(\zeta) \\ &\le C_1 C_2 \int_{\mathbb{T}} A_{q,\nu}(f)(\zeta) C_{q',\nu}(g)(\zeta)\, dm(\zeta) \\ &\lesssim \|f\|_{T^1_q(\nu)} \|g\|_{T^{\infty}_{q'}(\nu)}. \end{aligned} \tag{2.4}$$

Now let us prove (2.3). Fubini's theorem yields

$$\int_{\mathbb{T}} \left(\int_{\Gamma^{h(\zeta)}(\zeta)} k(z)\, d\nu(z) \right) dm(\zeta) = \int_{\mathbb{D}} |(I(z) \cap H(z)| k(z)\, d\nu(z),$$

where $H(z) = \{\zeta \in \mathbb{T} : \frac{1}{1+h(\zeta)} \le |z|\}$, so it suffices to show that

$$\frac{|(I(z) \cap H(z)|}{|I(z)|} \ge \frac{1}{C_2} \tag{2.5}$$

for all $z \in \mathbb{D}$. We will prove this only for z close enough to the boundary $\mathbb{T}$, the proof for other values of z follows from this reasoning with appropriate modifications. For $|z| \ge 1 - \frac{1}{n}$, set $z' = (1 - n(1 - |z|))z/|z|$ and $x = \frac{1}{|z|} -$

1, where n is a natural number ≥ 2 chosen such that $I(z) \cap I(u) = \emptyset$ if $u \notin T(z') \bigcup \left(\mathbb{D} \setminus \overline{D(0,|z|)}\right)$. This together with Fubini's theorem gives

$$\begin{aligned}
&\frac{1}{|I(z)|} \int_{I(z)} \left(\int_{\Gamma^x(\zeta)} |g(u)|^{q'} \, d\nu(u) \right) dm(\zeta) \\
&= \frac{1}{|I(z)|} \int_{\{|z|<|u|<1\}} |I(z) \cap I(u)| |g(u)|^{q'} \, d\nu(u) \\
&\leq \frac{1}{|I(z)|} \int_{T(z')} |I(z) \cap I(u)| |g(u)|^{q'} \, d\nu(u) \\
&\leq \frac{C_3}{|I(z')|} \int_{T(z')} |g(u)|^{q'} (1-|u|) \, d\nu(u) \leq C_3 \inf_{v \in I(z)} C^{q'}_{q',\nu}(g)(v),
\end{aligned} \tag{2.6}$$

where the last inequality is valid because

$$\frac{1}{|I(z')|} \int_{T(z')} |g(u)|^{q'} (1-|u|) \, d\nu(u) \leq \sup_{a \in \Gamma(v)} \frac{1}{|I(a)|} \int_{T(a)} |g(u)|^{q'} (1-|u|) \, d\nu(u)$$

for all $v \in I(z)$. Denote $E(z) = \mathbb{T} \setminus H(z) = \{\zeta \in \mathbb{T} : (1+h(\zeta))|z| < 1\}$. By the definition of $h(\zeta)$ and (2.6), and by choosing C_1 sufficiently large so that $C_1^{q'} > 2C_3$, we deduce

$$\begin{aligned}
|I(z) \cap E(z)| &\leq \int_{I(z)} \frac{A^{q'}_{q',\nu}(g|x)(\zeta)}{C_1^{q'} C^{q'}_{q',\nu}(g)(\zeta)} \, dm(\zeta) \\
&\leq \frac{1}{C_1^{q'} \inf_{v \in I(z)} C^{q'}_{q',\nu}(g)(v)} \int_{I(z)} A^{q'}_{q',\nu}(g|x)(\zeta) \, dm(\zeta) \\
&\leq \frac{C_3 |I(z)|}{C_1^{q'}} < \frac{1}{2} |I(z)|.
\end{aligned}$$

Therefore,

$$\frac{|I(z) \cap E(z)|}{|I(z)|} = 1 - \frac{|I(z) \cap E(z)|}{|I(z)|} \geq \frac{1}{2}, \quad |z| \geq 1 - \frac{1}{n},$$

and the inequality (2.5) follows. □

The reverse implication of Theorem A [11] can be proved by using geometric ideas, the boundedness of maximal functions and interpolation theorems on $L^p L^q(\nu, m)$ due to Benedek and Panzone [2].

The analogue of the following result on $\mathbb{R}^n \times (0,\infty)$ was proved in [11, Proposition 1]. See also [13, Lemma 4].

Lemma C. *Let $0 < p < \infty$ and let ν be a positive Borel measure on $\mathbb{D}$, finite on compact sets. Then there exists $\lambda_0 = \lambda_0(p) \geq 1$ such that*

$$\int_{\mathbb{T}} \left(\int_{\mathbb{D}} \left(\frac{1-|z|}{|1-\overline{\zeta} z|} \right)^{\lambda} d\nu(z) \right)^p dm(\zeta) \asymp \int_{\mathbb{T}} (\nu(\Gamma(\zeta)))^p \, dm(\zeta) + \nu(\{0\}) \tag{2.7}$$

for each $\lambda > \lambda_0$.

We defined the tent space $T_q^p(\nu)$ by using the lenses $\Gamma(\zeta)$. Different types of non-tangential approach regions could be used and they would induce the same spaces. In particular, the proof of Lemma C shows that we may replace $\Gamma(\zeta)$ by $\Gamma_\alpha(\zeta)$ for any $\alpha \in (0,\pi)$ in (2.7), and consequently the space $T_q^p(\nu)$ is independent of the aperture of the lens appearing in the definition, and the quasi-norms obtained for different lenses are equivalent.

Recall that $Z = \{z_k\}_{k=0}^\infty \subset \mathbb{D}$ is called a separated sequence if it is separated in the pseudohyperbolic metric, it is an ε-net if $\mathbb{D} = \bigcup_{k=0}^\infty \Delta(z_k,\varepsilon)$, and finally it is a δ-lattice if it is a 5δ-net and separated with constant $\gamma = \delta/5$. If $\nu = \sum_k \delta_{z_k}$, then we write $T_q^p(\nu) = T_q^p(\{z_k\})$. The next result [11, Theorem 2] (see also [13, Lemma 6]) plays an essential role in the proof of Theorem 1 (a).

Lemma D. *Let $0 < p < \infty$ and let $\{z_k\}$ be a separated sequence. Define*

$$S_\lambda(f)(z) = \sum_k f(z_k)\left(\frac{1-|z_k|}{1-\overline{z}_k z}\right)^\lambda, \quad z \in \mathbb{D}.$$

Then $S_\lambda : T_2^p(\{z_k\}) \to H^p$ is bounded for all $\lambda > \lambda_0$, where $\lambda_0 = \lambda_0(p) \geq 1$ is that of Lemma C.

We shall also use the following inequality. Here and on the following $\triangle$ denotes the Laplacian.

Lemma 3. *If $q \geq 2$ and $0 < r < 1$ there is a constant $C(q,r) > 0$ such that*

$$|f'(z)|^q(1-|z|^2)^q \leq C(q,r)\int_{\Delta(z,r)} \triangle|f|^q(\zeta)\,dA(\zeta), \quad z \in \mathbb{D}. \tag{2.8}$$

Proof. Let $r \in (0,1)$ be fixed. The classical Hardy-Stein-Spencer identity $\|f\|_{H^q}^q = |f(0)|^q + \frac{1}{2}\int_{\mathbb{D}} \triangle|f(z)|^q \log\frac{1}{|z|}\,dA(z)$ and the fact that the Laplacian $\triangle|f|^q$ is subharmonic when $q \geq 2$ give

$$\begin{aligned} |f'(0)|^q &\leq \left(\|f\|_{H^2}^2 - |f(0)|^2\right)^{\frac{q}{2}} \leq \|f\|_{H^2}^q - |f(0)|^q \leq \|f\|_{H^q}^q - |f(0)|^q \\ &= \frac{1}{2}\int_{\mathbb{D}} \triangle|f|^q(z)\log\frac{1}{|z|}\,dA(z) \leq C(q)\int_{\mathbb{D}} \triangle|f|^q(z)(1-|z|)\,dA(z). \end{aligned}$$

An application of this inequality to the function $f(rz)$ gives

$$|f'(0)|^q \leq C(q,r)\int_{\Delta(0,r)} \triangle|f|^q(z)\left(1-\frac{|z|}{r}\right)dA(z).$$

Replace now f by $f\circ\varphi_z$ to obtain

$$\begin{aligned} |f'(z)|^q(1-|z|^2)^q &\leq C(q,r)\int_{\Delta(z,r)} \triangle|f|^q(\zeta)\left(1-\frac{|\varphi_z(\zeta)|}{r}\right)dA(\zeta) \\ &\leq C(q,r)\int_{\Delta(z,r)} \triangle|f|^q(\zeta)\,dA(\zeta), \quad z \in \mathbb{D}. \end{aligned}$$

□

3. Proof of main results

We begin with proving Theorem 1(a).

Theorem 4. *Let* $0 < q \leq p < \infty$, $n \in \mathbb{N}$ *and let* μ *be a positive Borel measure on* $\mathbb{D}$. *Then* $D^{(n)} : H^p \to L^q(\mu)$ *is compact if and only if, for any fixed* $r \in (0,1)$, *the function*

$$\Phi_\mu(z) = \frac{\mu(\Delta(z,r))}{(1-|z|)^{1+qn}}$$

satisfies that

(i) $\Phi_\mu \in T^{\frac{p}{p-q}}_{\frac{2}{2-q}}(h)$, *if* $q < \min\{2,p\}$;

(ii) $\lim_{|a|\to 1^-} \frac{1}{|I(a)|}\int_{T(a)} |\Phi_\mu(z)|^{\frac{2}{2-p}} \frac{dA(z)}{1-|z|} = 0$, *if* $q = p < 2$;

(iii) $\lim_{R\to 1^-} \int_{\mathbb{T}} \left(\sup_{z\in\Gamma(\zeta)\setminus\overline{D(0,R)}} \Phi_\mu(z)\right)^{\frac{p}{p-q}} dm(\zeta) = 0$, *if* $2 \leq q < p$.

Proof. Recall the known estimate [9, Lemma 2.1]

$$|f^{(n)}(z)|^s \lesssim \frac{1}{(1-|z|)^{2+ns}} \int_{\Delta(z,r)} |f(\zeta)|^s \, dA(\zeta), \quad z \in \mathbb{D},\ s > 0. \tag{3.1}$$

Then, the above inequality and Fubini's theorem give

$$\begin{aligned}
\|f^{(n)}\|^q_{L^q(\mu)} &\lesssim \int_{\mathbb{D}} \frac{1}{(1-|z|)^{2+(n-1)q}} \int_{\Delta(z,r)} |f'(w)|^q \, dA(w)\, d\mu(z) \\
&\asymp \int_{\mathbb{D}} |f'(w)|^q \frac{\mu(\Delta(w,r))}{(1-|w|)^{3+(n-1)q}} |I(w)| \, dA(w) \\
&= \int_{\mathbb{D}} \left[|f'(w)|(1-|w|)\right]^q \Phi_\mu(w)|I(w)| \, dh(w)
\end{aligned} \tag{3.2}$$

Let $\{f_k\}_{k=1}^\infty$ such that $\sup_k ||f_k||_{H^p} < \infty$. Then, there is subsequence $\{f_{n_k}\}_{k=1}^\infty$ which converges uniformly on compact subsets of $\mathbb{D}$ to an analytic function f. Let denote $g_k = f_{n_k} - f$, $G_k(w) = |g'_k(w)|^q(1-|w|)^q$ and $dh_R = dh\chi_{\{R<|z|<1\}}$, $0 \leq R < 1$. Now, we shall show that conditions (i)-(iii) are sufficient.

(i). Fix $\varepsilon > 0$. Since $\Phi_\mu \in T^{\frac{p}{p-q}}_{\frac{2}{2-q}}(h)$, by the dominated convergence theorem there is R_0 such that

$$\sup_{R\geq R_0} ||\Phi_\mu||_{T^{\frac{p}{p-q}}_{\frac{2}{2-q}}(h_R)} < \varepsilon^q.$$

Next, choose k_0 with $|g_k(z)| < \varepsilon$ for any $k \geq k_0$ and $|z| \leq R_0$. Then, bearing in mind (3.2) and (2.1) and the inequality $\|G_k\|_{T^{\frac{p}{q}}_{\frac{2}{q}}(h)} \lesssim \|g_k\|^q_{H^p}$ (see (1.1))

$$\begin{aligned}
\|g_k^{(n)}\|^q_{L^q(\mu)} &\lesssim \varepsilon^q \int_{|w|\leq R_0} (1-|w|)^q \Phi_\mu(w)|I(w)|\,dh(w) \\
&\quad + \int_{\mathbb{D}} G_k(w)\Phi_\mu(w)|I(w)|\,dh_{R_0}(w) \\
&= \varepsilon^q \langle (1-|w|)^q, \Phi_\mu\rangle_{T^2_2(h)} + \langle G_k, \Phi_\mu\rangle_{T^2_2(h_{R_0})} \\
&\leq \varepsilon^q \|(1-|w)^q\|_{T^{\frac{p}{q}}_{\frac{2}{q}}(h)} \|\Phi_\mu\|_{T^{\left(\frac{p}{q}\right)'}_{\left(\frac{2}{q}\right)'}(h)} + \|G_k\|_{T^{\frac{p}{q}}_{\frac{2}{q}}(h)} \|\Phi_\mu\|_{T^{\left(\frac{p}{q}\right)'}_{\left(\frac{2}{q}\right)'}(h_{R_0})} \\
&\lesssim \varepsilon^q \left(\|\Phi_\mu\|_{T^{\left(\frac{p}{q}\right)'}_{\left(\frac{2}{q}\right)'}(h)} + ||g_k||^q_{H^p} \right) \lesssim \varepsilon^q,
\end{aligned}$$

So $D^{(n)} : H^p \to L^q(\mu)$ is compact. This together with [10, Theorem 1(i)] proves (i).

(ii). An standard argument (see [3, Theorem 3.4] for details) gives that $\lim_{|a|\to 1^-} \frac{1}{|I(a)|} \int_{T(a)} |\Phi_\mu(z)|^{\left(\frac{2}{p}\right)'} (1-|z|)\,dh(z) = 0$ if and only if

$$\lim_{R\to 1^-} \sup_{a\in\mathbb{D}} \frac{1}{|I(a)|} \int_{T(a)} |\Phi_\mu(z)|^{\left(\frac{2}{p}\right)'} (1-|z|)\,dh_R(z) = \lim_{R\to 1^-} \|\Phi_\mu\|_{T^\infty_{\left(\frac{2}{p}\right)'}(h_R)} = 0.$$

So fixed $\varepsilon > 0$, there is R_0 such that

$$\sup_{a\in\mathbb{D}, R\geq R_0} \frac{1}{|I(a)|} \int_{T(a)} |\Phi_\mu(z)|^{\left(\frac{2}{p}\right)'} (1-|z|)\,dh_R(z) = \sup_{R\geq R_0} \|\Phi_\mu\|_{T^\infty_{\left(\frac{2}{p}\right)'}(h_R)} < \varepsilon^p.$$

Let k_0 be such that $\sup_{k\geq k_0, |z|\leq R_0} |g_k(z)| < \varepsilon$. Then, by (3.2), (2.4) and (1.1)

$$\begin{aligned}
\|g_k^{(n)}\|^p_{L^p(\mu)} &\lesssim \varepsilon^p \langle (1-|w|)^p, \Phi_\mu\rangle_{T^2_2(h)} + \langle G_k, \Phi_\mu\rangle_{T^2_2(h_{R_0})} \\
&\lesssim \varepsilon^p \|(1-|w)^p\|_{T^1_{\frac{2}{p}}(h)} \|\Phi_\mu\|_{T^\infty_{\left(\frac{2}{p}\right)'}(h)} + \|G_k\|_{T^1_{\frac{2}{p}}}(h) \|\Phi_\mu\|_{T^\infty_{\left(\frac{2}{p}\right)'}(h_{R_0})} \\
&\lesssim \varepsilon^p (\|\Phi_\mu\|_{T^\infty_{\left(\frac{2}{p}\right)'}(h)} + ||g_k||^p_{H^p}) \lesssim \varepsilon^p,
\end{aligned}$$

which implies that $D^{(n)} : H^p \to L^p(\mu)$ is compact.

(iii). Let us observe that (2.8) and Fubini's theorem give

$$
\begin{aligned}
\||G_k\|^{\frac{q}{p}}_{T_1^{\frac{p}{q}}(h)} &= \int_{\mathbb{T}} \left(\int_{\Gamma(\zeta)} |g_k'(w)|^q (1-|w|)^q \, dh(w) \right)^{\frac{p}{q}} dm(\zeta) \\
&\lesssim \int_{\mathbb{T}} \left(\int_{\Gamma(\zeta)} \int_{\Delta(w,r)} \triangle |g_k|^q(z) \, dA(z) \, dh(w) \right)^{\frac{p}{q}} dm(\zeta) \\
&\leq \int_{\mathbb{T}} \left(\int_{\Gamma'(\zeta)} \triangle |g_k|^q(z) \, dA(z) \right)^{\frac{p}{q}} dm(\zeta)
\end{aligned}
$$

where $\Gamma'(\zeta) = \{z : \Gamma(\zeta) \cap \Delta(z,r) \neq \emptyset\}$. Using Lemma C and a result by Calderón [12, Theorem 1.3], we get $G_k \in T_1^{\frac{p}{q}}(h)$ with $\|G_k\|_{T_1^{\frac{p}{q}}(h)} \lesssim \|g_k\|^q_{H^p}$. From now on, the proof is analogous to both previous cases, so it will be omitted.

Reciprocally, assume that $D^{(n)} : H^p \to L^q(\mu)$ is compact. Let $\{z_k\}$ be a δ-lattice such that $z_k \neq 0$ for all k and let $\mathcal{C}T_2^p(\{z_k\}) = \{f \in T_2^p(\{z_k\}) : ||f||_{T_2^p(\{z_k\})} = 1\}$. For each $R \in [0,1)$ and $\lambda > \lambda_0$ (λ_0 is that of Lemma D) consider the operator

$$
S_{\lambda,\,R}(f)(z) = \sum_{|z_k| \geq R} f(z_k) \left(\frac{1-|z_k|}{1-\overline{z}_k z} \right)^{\lambda}, \quad z \in \mathbb{D}.
$$

Let us observe that $S_{\lambda,\,0}(f) = S_\lambda(f)$. By Lemma D, there exists $C > 0$ such that

$$
||S_{\lambda,\,R}(f)||_{H^p} \leq C ||f||_{T_2^p(\{z_k\})}, \quad \text{for each } R \in [0,1) \ .
$$

So by the assumption the closure of the set $\left\{ D^{(n)} \circ S_{\lambda,\,R} \left(\mathcal{C}T_2^p(\{z_k\}) \right) \right\}_{R \in [0,1)}$ is compact in $L^q(\mu)$. So, fixed $\varepsilon > 0$, standard arguments assert that there is ρ such that

(3.3)

$$
\int_{\rho<|z|<1} |D^{(n)} \circ S_{\lambda,\,R}(f)(z)|^q \, d\mu(z) < \varepsilon^q \quad \text{for any } R \in [0,1) \text{ and } f \in \mathcal{C}T_2^p(\{z_k\}).
$$

Since $\{z_k\}$ is separated and $\lambda > 1$, there is R_0 such that for any $R \geq R_0$ $\sum_{|z_k|\geq R}(1-|z_k|)^\lambda < \varepsilon^2$. Joining this with Lemma C, we get

$$\begin{aligned}
|D^{(n)} \circ S_{\lambda, R}(f)(z)| &\leq C(\rho, n) \sum_{|z_k|\geq R} |f(z_k)|(1-|z_k|)^\lambda \\
&\leq C_\rho \left(\sum_{|z_k|\geq R} |f(z_k)|^2 (1-|z_k|)^\lambda \right)^{\frac{1}{2}} \varepsilon \\
&\leq C(\rho, n)\varepsilon \inf_{\zeta\in\mathbb{T}} \left(\sum_k |f(z_k)|^2 \left| \frac{1-|z_k|}{1-\overline{z}_k\zeta} \right|^\lambda \right)^{\frac{1}{2}} \\
&\leq C(\rho, n)\varepsilon \int_{\mathbb{T}} \left(\sum_k |f(z_k)|^2 \left(\frac{1-|z_k|}{|1-\overline{z}_k\zeta|} \right)^\lambda \right)^{\frac{p}{2}} dm(\zeta) \\
&\leq C(\rho, n)\varepsilon \|f\|^p_{T^p_2(\{z_k\})}, \quad \text{for any } |z| \leq \rho \text{ and } R \geq R_0,
\end{aligned}$$

which together with (3.3) gives that

$$||D^{(n)} \circ S_{\lambda, R}(f)||_{L^q(\mu)} \lesssim \varepsilon \|f\|_{T^p_2(\{z_k\})}, \quad \text{for all } R \geq R_0 \text{ and } f \in T^p_2(\{z_k\}). \tag{3.4}$$

That is

$$\int_{\mathbb{D}} \left| \sum_{|z_k|\geq R} f(z_k) \frac{(1-|z_k|)^\lambda}{(1-\overline{z}_k z)^{\lambda+n}} \right|^q d\mu(z) \lesssim \varepsilon^q \|f\|^p_{T^q_2(\{z_k\})},$$

for all $R \geq R_0$ and $f \in T^p_2(\{z_k\})$. Replace now $f(z_k)$ by $f(z_k)r_k(t)$, where r_k denotes the kth Rademacher function, and integrate with respect to t to obtain

$$\int_0^1 \int_{\mathbb{D}} \left| \sum_{|z_k|\geq R} f(z_k) \frac{(1-|z_k|)^\lambda}{(1-\overline{z}_k z)^{\lambda+n}} r_k(t) \right|^q d\mu(z)\, dt \lesssim \varepsilon^q \|f\|^q_{T^p_2(\{z_k\})},$$

from which Fubini's theorem and an application of Khinchine's inequality yield

$$I = \int_{\mathbb{D}} \left(\sum_{|z_k|\geq R} |f(z_k)|^2 \frac{(1-|z_k|)^{2\lambda}}{|1-\overline{z}_k z|^{2\lambda+2n}} \right)^{\frac{q}{2}} d\mu(z) \lesssim \varepsilon^q \|f\|^q_{T^p_2(\{z_k\})}.$$

Now, for any fixed $r \in (0,1)$,

$$\begin{aligned}
I &\gtrsim \sum_{|z_j|\geq R} \int_{\Delta(z_j, r)} \left(\sum_{|z_k|\geq R} |f(z_k)|^2 \frac{(1-|z_k|)^{2\lambda}}{|1-\overline{z}_k z|^{2\lambda+2n}} \right)^{\frac{q}{2}} d\mu(z) \\
&\gtrsim \sum_{|z_j|\geq R} \left(|f(z_j)|^2 \frac{(1-|z_j|)^{2\lambda}}{(1-|z_j|)^{2\lambda+2n}} \right)^{\frac{q}{2}} \mu(\Delta(z_j, r)) \\
&= \sum_{|z_j|\geq R} \frac{|f(z_j)|^q}{(1-|z_j|)^{qn}} \mu(\Delta(z_j, r)),
\end{aligned}$$

and hence

$$
\begin{aligned}
&\sum_{|z_j|\geq R} \frac{|f(z_j)|^q}{(1-|z_j|)^{qn}} \mu(\Delta(z_j,r)) \\
&= \sum_{|z_j|\geq R} |f(z_j)|^q \left(\frac{\mu(\Delta(z_j,r))}{(1-|z_j|)^{1+qn}}\right)(1-|z_j|) \lesssim \varepsilon^q \|f\|^q_{T_2^p(\{z_k\})} \\
&= \varepsilon^q \left(\int_{\mathbb{T}} \left(\sum_{z_k\in\Gamma(\zeta)} |f(z_k)|^2\right)^{\frac{p}{2}} dm(\zeta)\right)^{\frac{q}{p}} \qquad (3.5) \\
&= \varepsilon^q \left(\int_{\mathbb{T}} \left(\left(\sum_{z_k\in\Gamma(\zeta)} (|f(z_k)|^q)^{\frac{2}{q}}\right)^{\frac{q}{2}}\right)^{\frac{p}{q}} dm(\zeta)\right)^{\frac{q}{p}}.
\end{aligned}
$$

(ii) If $q = p < 2$, then $s = \frac{p}{q} = 1$ and $v = 2/p > 1$, so by Theorem A $\left(T_v^1(\{z_k\})\right)^\star \simeq T_{v'}^\infty(\{z_k\})$ with equivalence of norms. Therefore (3.5) yields

$$
\sup_{a\in\mathbb{D}} \frac{1}{|I(a)|} \sum_{z_k\in T(a),|z_k|\geq R} \left(\frac{\mu(\Delta(z_k,r))}{(1-|z_k|)^{1+pn}}\right)^{\frac{2}{2-p}} |I(z_k)| \lesssim \varepsilon^p
$$

for all $R \geq R_0$. The above inequality is a discrete version of,

$$
\sup_{a\in\mathbb{D},R\geq R_0} \frac{1}{|I(a)|} \int_{T(a)\cap\{R<|z|<1\}} |\Phi_\mu(z)|^{\left(\frac{2}{p}\right)'} \frac{dA(z)}{1-|z|} \lesssim \varepsilon^p.
$$

So, $\lim_{R\to 1^-} \left(\sup_{a\in\mathbb{D}} \frac{1}{|I(a)|} \int_{T(a)\cap\{R<|z|<1\}} |\Phi_\mu(z)|^{\left(\frac{2}{p}\right)'} \frac{dA(z)}{1-|z|}\right) = 0$, which is equivalent to $\lim_{|a|\to 1^-} \frac{1}{|I(a)|} \int_{T(a)} |\Phi_\mu(z)|^{\left(\frac{2}{p}\right)'} \frac{dA(z)}{1-|z|} = 0$.

(iii) If $2 < q < p$, then $s = \frac{p}{q} > 1$ and $v = \frac{2}{q} < 1$, and hence [10, Proposition 3] yields

$$
\int_{\mathbb{T}} \left(\sup_{z_k\in\Gamma(\zeta)\setminus\overline{D(0,R)}} \frac{\mu(\Delta(z_k,r))}{(1-|z_k|)^{1+qn}}\right)^{\frac{p}{p-q}} dm(\zeta) < \varepsilon^q, \quad \text{for all } R \geq R_0.
$$

from which the assertion follows. The case $q = 2$ is proved similarly by using Theorem A instead of [10, Proposition 3]. This finishes the proof. □

Now we deal with the second part of Theorem 1(b).

Theorem 5. *Let either $0 < p < q < \infty$ or $2 \leq p = q < \infty$ and $n \in \mathbb{N}$, and let μ be a positive Borel measure on $\mathbb{D}$. Then the following conditions are equivalent:*

(i) *$D^{(n)} : H^p \to L^q(\mu)$ is compact;*

(ii) *$\lim_{|z|\to 1^-} \frac{\mu(S(z))}{(1-|z|)^{nq+\frac{q}{p}}} = 0$;*

(iii) *$\lim_{|z|\to 1^-} \frac{\mu(\Delta(z,r))}{(1-|z|)^{nq+\frac{q}{p}}} = 0$ for any fixed $r \in (0,1)$.*

Proof. A proof of (i)$\Rightarrow$(ii) (and (i)$\Rightarrow$(iii)) can be obtained by using the test functions $f_a(z)=\left(\frac{1-|a|^2}{(1-\overline{a}z)^2}\right)^{1/p}$, $a\in\mathbb{D}$, and a regular reasoning. So it is omitted.

It is enough to prove (iii)$\Rightarrow$(i). We shall split the proof in two cases.

Case $\mathbf{0<p<q<\infty}$. The argument follows ideas from the proof of [9, Theorem 3.1]. Let $r\in(0,1)$ be fixed, choose $s\in(p,q)$ and denote

$$d\mu^\star(\zeta)=(1-|\zeta|)^{\frac{s}{p}-2}\,dA(\zeta),\quad \zeta\in\mathbb{D}.$$

Let $\{f_k\}_{k=1}^\infty$ such that $\sup_k ||f_k||_{H^p}<\infty$. Then, there is subsequence $\{f_{n_k}\}_{k=1}^\infty$ which converges uniformly on compact subsets of $\mathbb{D}$ to to an analytic function f. Let denote $g_k=f_{n_k}-f$. Fix $\varepsilon>0$, by hypothesis there is ρ such that

$$\frac{\mu\left(\Delta(z,r)\right)}{(1-|z|)^{nq+\frac{q}{p}}}<\varepsilon^q,\quad \text{if } \rho<|z|<1.$$

On the other hand, there is k_0 such that $|g_k(z)|<\varepsilon$ for any $k\ge k_0$ and $|z|\le\rho$. So, bearing in mind (3.1) and Minkowski's inequality in continuous form

$$\begin{aligned}\|g_k^{(n)}\|^q_{L^q(\mu)} &\lesssim \int_{\mathbb{D}}\left(\frac{1}{(1-|z|)^{2+ns}}\int_{\Delta(z,r)}|g_k(\zeta)|^s\,dA(\zeta)\right)^{\frac{q}{s}}d\mu(z)\\ &\lesssim \left(\int_{\mathbb{D}}|g_k(\zeta)|^s\frac{(\mu(\Delta(\zeta,r)))^{\frac{s}{q}}}{(1-|\zeta|)^{ns}}\,dh(\zeta)\right)^{\frac{q}{s}}\\ &\lesssim \varepsilon^q\left(\int_{|\zeta|\le\rho}\frac{(\mu(\Delta(\zeta,r)))^{\frac{s}{q}}}{(1-|\zeta|)^{ns}}\,dh(\zeta)\right)^{\frac{q}{s}}+\varepsilon^q\left(\int_{\rho<|\zeta|<1}|g_k(\zeta)|^s\,d\mu^\star(\zeta)\right)^{\frac{q}{s}}\\ &\lesssim \varepsilon^q\left(\mu^\star(\mathbb{D})^{\frac{q}{s}}+||g_k||^q_{L^s(\mu^\star)}\right).\end{aligned}$$

Next, since $\mu^\star(S(a))\lesssim(1-|a|)^{\frac{s}{p}}$ for $a\in\mathbb{D}$, by [7, Theorem 9.4] $||g_k||_{L^s(\mu^\star)}\lesssim ||g_k||_{H^p}$, which together with the above inequalities implies that $D^{(n)}:H^p\to L^q(\mu)$ is compact.

Case $\mathbf{q=p\ge 2}$. By (3.1), (2.8) and Fubini's theorem

$$\begin{aligned}\|f^{(n)}\|^p_{L^p(\mu)} &\lesssim \int_{\mathbb{D}}\frac{1}{(1-|z|)^{2+(n-1)p}}\left(\int_{\Delta(z,\rho)}|f'(\zeta)|^p\,dA(\zeta)\right)d\mu(z)\\ &\lesssim \int_{\mathbb{D}}\frac{1}{(1-|z|)^{2+np}}\left(\int_{\Delta(z,\rho)}\left(\int_{\Delta(\zeta,s)}\triangle|f|^p(u)\,dA(u)\right)dA(\zeta)\right)d\mu(z)\\ &\lesssim \int_{\mathbb{D}}\frac{1}{(1-|z|)^{np}}\left(\int_{\Delta(z,r)}\triangle|f|^p(u)\,dA(u)\right)d\mu(z)\\ &\asymp \int_{\mathbb{D}}\frac{\triangle|f|^p(u)}{(1-|u|)^{np}}\mu(\Delta(u,r))\,dA(u),\end{aligned}$$

where $\rho, s \in (0,1)$ are chosen sufficiently small depending only on r. Putting together this inequality with the Hardy-Stein-Spencer identity $\|f\|_{H^p}^p = |f(0)|^p + \frac{1}{2}\int_{\mathbb{D}} \triangle |f(z)|^p \log \frac{1}{|z|}\, dA(z)$, the proof can be finished as in the previous case.

□

The following result contains Theorem 2.

Theorem 6. *Let $0 < q < p < \infty$ and let μ be a positive Borel measure on $\mathbb{D}$. Then the following conditions are equivalent:*

- (i) $I_d : H^p \to L^q(\mu)$ *is compact;*
- (ii) $I_d : H^p \to L^q(\mu)$ *is bounded;*
- (iii) *The function*
$$\Psi_\mu(\zeta) = \int_{\mathbb{D}} \left(\frac{1-|z|}{|1-\overline{\zeta}z|} \right)^\lambda \frac{d\mu(z)}{1-|z|}$$
belongs to $L^{\frac{p}{p-q}}(\mathbb{T}, m)$ *for all* $\lambda > 0$ *large enough;*
- (iv) *The function* $B_\mu(\zeta) = \int_{\Gamma(\zeta)} \frac{d\mu(z)}{1-|z|}$ *belongs to* $L^{\frac{p}{p-q}}(\mathbb{T}, m)$*;*
- (v) *For each* $0 < r < 1$*, the function*
$$\zeta \mapsto \int_{\Gamma(\zeta)} \frac{\mu\left(\Delta(z,r)\right)}{(1-|z|)^3}\, dA(z)$$
belongs to $L^{\frac{p}{p-q}}(\mathbb{T}, m)$*;*
- (vi) $M(\mu)(z) = \sup_{z \in S(a)} \frac{\mu(S(a))}{1-|a|} \in L^{\frac{p}{p-q}}(\mathbb{T}, m)$.

Proof. The equivalences $(ii) \Leftrightarrow (iii) \Leftrightarrow (iv) \Leftrightarrow (vi)$ follows from [10, Section D]. Next, fixed $0 < r < 1$, let us observe that for a positive measure ν

(3.6)
$$\begin{aligned}\int_{\mathbb{D}} \left(\frac{1-|z|}{|1-\overline{\zeta}z|} \right)^\lambda d\nu(z) &= \int_{\mathbb{D}} \left(\int_{\Delta(u,r)} \left(\frac{1-|z|}{|1-\overline{\zeta}z|} \right)^\lambda \frac{1}{|\Delta(z,r)|} d\nu(z) \right) dA(u) \\ &\asymp \int_{\mathbb{D}} \left(\frac{1-|u|}{|1-\overline{\zeta}u|} \right)^\lambda \nu\left(\Delta(u,r)\right)\, dh(u), \quad \text{for all } \zeta \in \mathbb{T}.\end{aligned}$$

So choosing $d\nu(z) = \frac{d\mu(z)}{1-|z|}$ and applying Lemma C, we have that $(iv) \Leftrightarrow (v)$.

Finally, let us see $(iv) \Rightarrow (i)$. By hypothesis and (3.6)

$$\int_{\mathbb{T}} \left(\int_{\mathbb{D}} \left(\frac{1-|u|}{|1-\overline{\zeta}u|} \right)^\lambda \frac{\mu\left(\Delta(u,r)\right)}{(1-|u|)^3}\, dA(u) \right)^{\frac{p}{p-q}} dm(\zeta) < \infty.$$

Then, by dominated convergence theorem and (3.6),

$$\begin{aligned} 0 &= \lim_{R \to 1^-} \int_{\mathbb{T}} \left(\int_{\{R<|z|<1\}} \left(\frac{1-|z|}{|1-\overline{\zeta}z|} \right)^\lambda \frac{d\mu(z)}{1-|z|} \right)^{\frac{p}{p-q}} dm(\zeta) \\ &\gtrsim \lim_{R \to 1^-} \int_{\mathbb{T}} \left(\int_{\Gamma(\zeta) \setminus \overline{D(0,R)}} \frac{d\mu(z)}{1-|z|} \right)^{\frac{p}{p-q}} dm(\zeta). \end{aligned}$$

This together with the equivalence $||f||_{H^p}^p \asymp \int_{\mathbb{T}} \left(\sup_{z\in\Gamma(\zeta)} |f(z)|\right)^p \, dm(\zeta)$ [8] and standard arguments, yields that $I_d : H^p \to L^q(\mu)$ is compact. □

References

[1] P. Ahern and J. Bruna, Maximal and area integral characterizations of Hardy-Sobolev spaces in the unit ball of $\mathbb{C}^n$, Rev. Mat. Iberoamericana 4 (1988), no. 1, 123–153.

[2] A. Benedek and R. Panzone, The spaces L^p, with mixed norm, Duke Math. J. 28 (1961), 301–324.

[3] O. Blasco and H. Jarchow, A note on Carleson measures for Hardy spaces, Acta Sci. Math. (Szeged) 71 (2005), no. 1-2, 371–389.

[4] W. S. Cohn and I. E. Verbitsky, Factorization of tent spaces and Hankel operators, J. Funct. Anal. 175 (2000), no. 2, 308–329.

[5] W. S. Cohn, S. H. Ferguson and R. Rochberg, Boundedness of higher order Hankel forms, Factorization in potential spaces and derivations, Proc. London Math. Soc. 82 (2001), no. 1, 110–130.

[6] R. R. Coifman, Y. Meyer and E. M. Stein, Some new functions spaces and their applications to Harmonic Analysis, J. Funct. Anal. 62 (1985), no. 3, 304–335.

[7] P. Duren, Theory of H^p Spaces, Academic Press, New York-London 1970.

[8] C. Fefferman and E. M. Stein, H^p spaces of several variables, Acta Math. 129 (1972), no. 3-4, 137-193.

[9] D. H. Luecking, Forward and reverse inequalities for functions in Bergman spaces and their derivatives, Amer. J. Math. 107, (1985), 85–111.

[10] D. H. Luecking, Embedding derivatives of Hardy spaces into Lebesgue spaces, Proc. London Math. Soc. 63 (1991), no. 3, 595–619.

[11] D. H. Luecking, Embedding theorems for spaces of analytic functions via Khinchine's inequality, Michigan Math. J. 40 (1993), no. 2, 333–358.

[12] M. Pavlović, On the Littlewood-Paley g-function and Calderon's area theorem, Expo. Math. 1 (2013), n. 31, 169-195.

[13] J. A. Peláez and J. Rättyä, Embedding theorems for Bergman spaces via harmonic analysis, to appear in Math. Annalen, DOI: 10. 1007/s00208-014-1108-5.

[14] K. Zhu and R. Zhao, Theory of Bergman spaces in the unit ball, Memoires de la SFM 115 (2008).

Departamento de Analisis Matematico, Universidad de Malaga, Campus de Teatinos, 29071 Malaga, Spain

E-mail address: `japelaez@uma.es`